arXiv:1406.7835v5 [math.NT] 1 Nov 2014

ON THE GAUSS *EΥPHKA* THEOREM AND SOME ALLIED INEQUALITIES

ALEXANDER BERKOVICH

Abstract. We use the 1907 Hurwitz formula along with the Jacobi triple product identity to understand representation properties of two JP (Jones-Pall) forms of Kaplansky: $9x^2+16y^2+36z^2+16yz+4xz+8xy$ and $9x^2+17y^2+32z^2-8yz+8xz+6xy$. We also discuss three nontrivial analogues of the Gauss *EΥPHKA* theorem. The technique used can be applied to all known spinor regular ternary quadratic forms.

1. Introduction

We use notation (a,b,c,d,e,f) to represent a positive ternary quadratic form $ax^2+by^2+cz^2+dyz+exz+fxy$. We remark that this paper only considers positive ternary quadratic forms. We use $(a,b,c,d,e,f;n)$ to denote the total number of representations of n by (a,b,c,d,e,f), and take $(a,b,c,d,e,f;n)=0$ when $n\notin\mathbb{N}$. The associated theta series to the form (a,b,c,d,e,f) is

$$\vartheta(a,b,c,d,e,f,q):=\sum_{x,y,z} q^{ax^2+by^2+cz^2+dyz+exz+fxy}=\sum_{n\geq 0}(a,b,c,d,e,f;n)q^n. \tag{1.1}$$

Here and throughout the paper we assume that q is a complex number with $|q|<1$. The discriminant Δ of (a,b,c,d,e,f) is defined as

$$\Delta:=\frac{1}{2}\det\begin{pmatrix} 2a & f & e\\ f & 2b & d\\ e & d & 2c\end{pmatrix}=4abc+def-ad^2-be^2-cf^2.$$

Note that the discriminant $\Delta>0$ for a positive ternary quadratic form. Two ternary quadratic forms of discriminant Δ are in the same genus if they are equivalent over $\mathbb{Q}$ via a transformation matrix in $SL(3,\mathbb{Q})$ whose entries have denominators coprime to 2Δ.

The following result is used more often than any other in research in the theory of numbers.

Theorem 1.1 (Three-square theorem)**.** *Let n,m and v be non-negative integers. Then*

$$(1,1,1,0,0,0;n)\geq 0, \tag{1.2}$$

with equality if and only if $n=4^v(8m+7)$.

The proof was given by Legendre in [19] and by Gauss in [11]. Using this result, Gauss proved his celebrated *EΥPHKA* theorem. According to this theorem, every natural number is a sum of three triangular numbers. This is equivalent to the statement that every natural number of the form $8m+3$ is a sum of three odd squares.

Theorem 1.2 (Gauss)**.** *Let n be a non-negative integer. Then*

$$(1,1,1,0,0,0;8n+3)=[q^{8n+3}]\sum_{\substack{x\equiv 1 \pmod 2,\\ y\equiv 1 \pmod 2,\\ z\equiv 1 \pmod 2}} q^{x^2+y^2+z^2}>0. \tag{1.3}$$

Date: November 4, 2014.

2010 *Mathematics Subject Classification.* 11B65, 11E12, 11E16, 11E20, 11E25, 11E41, 11F37.

Key words and phrases. θ-function identities, representation properties, spinor regular ternary quadratic forms, 1907 Hurwitz formula.

Here and throughout the paper $[q^n]F(q)$ denotes the coefficient of q^n extracted from $F(q)$ (when written as a Maclaurin series).

Using the Three-square theorem along with two easily verifiable identities

$$(1,1,2,0,0,0;2n)=(1,1,1,0,0,0;n), n\in\mathbb{N}, \tag{1.4}$$

$$3(1,1,2,0,0,0;2n+1)=(1,1,1,0,0,0;4n+2), n\in\mathbb{N}, \tag{1.5}$$

we can prove the following:

Theorem 1.3. *Let n, m and v be non-negative integers. Then*

$$(1,1,2,0,0,0;n)\geq 0, \tag{1.6}$$

with equality if and only if $n=4^v2(8m+7)$.

This result is, essentially, Dickson's Theorem 86 in [10]. Theorem 89 in [10] can be stated as

Theorem 1.4. *Let n, m and v be non-negative integers. Then*

$$(1,1,3,0,0,0;n)\geq 0, \tag{1.7}$$

with equality if and only if $n=9^v3(3m+2)$.

In his 1907 letter, Hurwitz [14] proposed that $(1,1,1,0,0,0;n^2)$ may be expressed as a simple finite function of the divisors of n. In fact, if n is a positive integer with prime factorization

$$n=2^a\prod_{p>2}p^v. \tag{1.8}$$

Then

$$\frac{(1,1,1,0,0,0;n^2)}{6}=\prod_{p>2}\left(\frac{1-p^{v+1}}{1-p}-\left(\frac{-1}{p}\right)\frac{1-p^v}{1-p}\right). \tag{1.9}$$

Here and throughout the paper $\left(\frac{a}{n}\right)$ denotes the Jacobi symbol. A proof of this formula, using integral quaternions, has been given by Pall [21]. A completely elementary arithmetical proof, based on the methods of Liouville, was given by Olds [20]. Recently, explicit analogues of Hurwitz's formula for many diagonal ternary quadratic forms of class number one were established in [9] and [12].

Theorem 1.5. *Let n be a positive integer with prime factorization*

$$n=2^a\prod_{p>2}p^v. \tag{1.10}$$

Then

$$\frac{(1,1,2,0,0,0;n^2)}{4}=f(a)\prod_{p>2}\left(\frac{1-p^{v+1}}{1-p}-\left(\frac{-2}{p}\right)\frac{1-p^v}{1-p}\right), \tag{1.11}$$

where $f(0)=1$ and $f(a)=3$ when $a\geq 1$.

Theorem 1.6. *Let n be a positive integer with prime factorization*

$$n=2^a3^b\prod_{p>3}p^v. \tag{1.12}$$

Then

$$\frac{(1,1,3,0,0,0;n^2)}{4}=(2^{a+1}-1)\prod_{p>3}\left(\frac{1-p^{v+1}}{1-p}-\left(\frac{-3}{p}\right)\frac{1-p^v}{1-p}\right). \tag{1.13}$$

We remark that all analogues of Hurwitz's formula for ternary quadratic forms of class number one easily follow from the Siegel formula [23], [4, (2.1)]. Moreover, it is not necessary for the class number to be one. For example, it is not difficult to establish the following:

Theorem 1.7. *Let n be a positive integer with prime factorization*

$$n = 2^a 3^b \prod_{p>3} p^v. \tag{1.14}$$

Then

$$\frac{(1,3,36,0,0,0;n^2)+(3,4,9,0,0,0;n^2)}{2} = (3\ 2^a-2)g(b)\prod_{p>3}\left(\frac{1-p^{v+1}}{1-p}-\left(\frac{-3}{p}\right)\frac{1-p^v}{1-p}\right), \tag{1.15}$$

where $g(0)=1, g(b)=2$ when $b\geq 1$.

Next, we observe that (1.9), (1.11), (1.13) and (1.15) imply the following inequalities:

Corollary 1.8. *Let $M \not\equiv 0 \pmod 2, E \not\equiv 0 \pmod 2$ and $W \not\equiv 0 \pmod 3$ be positive integers. Then*

$$\frac{(1,1,1,0,0,0;M^2)}{6} \geq M, \tag{1.16}$$

with equality if and only if all prime divisors of M are congruent to $1 \pmod 4$,

$$\frac{(1,1,2,0,0,0;E^2)}{4} \geq E, \tag{1.17}$$

with equality if and only if all prime divisors of E are congruent to $1 \pmod 8$ or $3 \pmod 8$,

$$\frac{(1,1,3,0,0,0;W^2)}{4} \geq W, \tag{1.18}$$

$$\frac{(1,3,36,0,0,0;W^2)+(3,4,9,0,0,0;W^2)}{2} \geq W, \tag{1.19}$$

with equality if and only if all prime divisors of W are congruent to $1 \pmod 3$.

One of main objectives of this paper is to discuss three nontrivial analogues of the Gauss theorem 1.2. We will use (1.16) to prove:

Theorem 1.9. *Let n be a positive integer. Then*

$$[q^{8n+1}] \sum_{\substack{x\equiv 1 \pmod 4,\\ y\equiv 2 \pmod 8,\\ z\equiv 2 \pmod 8}} q^{x^2+y^2+z^2} \geq 0, \tag{1.20}$$

with equality if and only if $8n+1=M^2$ and all prime divisors of M are congruent to 1 modulo 4.

We will use (1.17) to prove:

Theorem 1.10. *Let n be a positive integer. Then*

$$[q^{8n+1}] \sum_{\substack{x\equiv 1 \pmod 4,\\ y\equiv 4 \pmod{16},\\ z\equiv 0 \pmod 2}} q^{x^2+y^2+2z^2} \geq 0, \tag{1.21}$$

with equality if and only if $8n+1= E^2$ and all prime divisors of E are congruent to 1 or 3 modulo 8.

We will use (1.18) to prove:

Theorem 1.11. *Let n be a positive integer. Then*

$$[q^{24n+1}] \sum_{\substack{x\equiv 3 \pmod{12},\\ y\equiv z \pmod 6,\\ y+z\equiv 2 \pmod 6}} q^{x^2+4y^2+12z^2} \geq 0, \tag{1.22}$$

with equality if and only if $24n+1=W^2$ and all prime divisors of W are congruent to 1 modulo 3.

The rest of this paper is organized as follows. In the next section, we collect necessary q-series identities. In Section 3 we will prove Theorems 1.9, 1.10 and 1.11. In Section 4, we prove some conjectures of Kaplansky regarding two spinor-regular positive ternary forms of discriminant 16384.

2. q-Series and Theta Function identities

We begin by collecting some standard q-notations, definitions, and useful formulas,

$$(a;q)_\infty := \prod_{n=0}^{\infty}(1-aq^n), \tag{2.1}$$

and

$$E(q) := (q;q)_\infty. \tag{2.2}$$

Note that

$$E(-q) := \frac{E^3(q^2)}{E(q^4)E(q)}. \tag{2.3}$$

Next, we recall the Ramanujan theta function

$$f(a,b) := \sum_{n=-\infty}^{\infty} a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}}, \qquad |ab|<1. \tag{2.4}$$

In Ramanujan's notation, the celebrated Jacobi triple product identity takes the shape [6, p. 35],

$$f(a,b) = (-a;ab)_\infty(-b;ab)_\infty(ab;ab)_\infty. \tag{2.5}$$

One may use (2.5) to derive the following important special cases:

$$f(-q,-q^2) = \sum_{n=-\infty}^{\infty}(-1)^n q^{\frac{n(3n+1)}{2}} = E(q), \tag{2.6}$$

$$\sum_{n>0}\left(\frac{-4}{n}\right) nq^{n^2} = qE(q^8)^3, \tag{2.7}$$

$$\sum_{n>0}\left(\frac{-2}{n}\right) nq^{n^2} = qE(-q^8)^3, \tag{2.8}$$

$$\phi(q) := f(q,q) = \sum_{n=-\infty}^{\infty} q^{n^2} = \frac{E^5(q^2)}{E^2(q^4)E^2(q)}, \tag{2.9}$$

$$\phi(-q) = \sum_{n=-\infty}^{\infty}(-1)^n q^{n^2} = \frac{E^2(q)}{E(q^2)}, \tag{2.10}$$

$$\psi(q) := f(q,q^3) = \sum_{n=-\infty}^{\infty} q^{2n^2-n} = \frac{E^2(q^2)}{E(q)}, \tag{2.11}$$

$$\psi(-q) = \sum_{n=-\infty}^{\infty}(-1)^n q^{2n^2-n} = \frac{E(q^4)E(q)}{E(q^2)}, \tag{2.12}$$

$$f(q,q^2) = \sum_{n=-\infty}^{\infty} q^{\frac{n(3n+1)}{2}} = \frac{E^2(q^3)E(q^2)}{E(q^6)E(q)}, \tag{2.13}$$

$$f(q,q^5) = \sum_{n=-\infty}^{\infty} q^{3n^2+2n} = \frac{E(q^{12})E(q^3)E^2(q^2)}{E(q^6)E(q^4)E(q)}. \tag{2.14}$$

Note that (2.6) is the famous Euler pentagonal number theorem. We observe that (2.10) and (2.11) can be combined to yield

$$\phi(-q^8)^2\psi(q^8) = E(q^8)^3. \tag{2.15}$$

Analogously, from (2.9), (2.10), (2.11) and (2.12), we deduce that

$$\phi(-q^2)\psi(q) = \phi(q)\psi(-q). \tag{2.16}$$

The relations (2.15) and (2.16) will come in handy in our proof of Theorems 1.9 and 1.10, respectively.

The well-known quintuple product identity can be formulated as in [7, p. 18],

$$\sum_{n=-\infty}^{\infty} q^{3n^2+n}\left(\frac{z^{3n}}{q^{3n}} - \frac{q^{3n+1}}{z^{3n+1}}\right) = (q^2;q^2)_\infty (qz;q^2)_\infty \left(\frac{q}{z};q^2\right)_\infty (z^2;q^4)_\infty \left(\frac{q^4}{z^2};q^4\right)_\infty, \tag{2.17}$$

with $z \neq 0$. It implies that

$$\sum_{n>0}\left(\frac{-12}{n}\right) nq^{n^2} = q\phi(q^{12})E(q^{12})^2 = q\frac{E(q^{24})^5}{E(q^{48})^2}, \tag{2.18}$$

and

$$\sum_{n>0}(-1)^{n+1}\left(\frac{-3}{n}\right) nq^{n^2} = q\phi(q^3)E(q^{12})^2 = q\frac{E(q^6)^5}{E(q^3)^2}. \tag{2.19}$$

The function $f(a,b)$ may be dissected in many different ways. We will use the following trivial dissections [6, p. 40, p. 49],

$$\phi(q) = \phi(q^4) + 2q\psi(q^8), \tag{2.20}$$

$$\phi(q) = \phi(q^9) + 2qf(q^3, q^{15}), \tag{2.21}$$

$$f(q, q^5) = f(q^8, q^{16}) + qf(q^4, q^{20}). \tag{2.22}$$

Next, we define

$$a(q) := \sum_{x,y} q^{x^2+xy+y^2} \tag{2.23}$$

and

$$c(q) := \sum_{x,y} q^{x^2+xy+y^2+x+y}. \tag{2.24}$$

These functions were extensively studied in recent literature [1], [6], [13] and [22]. I will record below some useful formulas from [3], [6] and [13]:

$$\phi(q)^2 = \phi(q^2)^2 + 4q\psi(q^4)^2, \tag{2.25}$$

$$\phi(q)^4 - \phi(q^3)^4 = 8qf(q, q^5)^3\phi(q^3), \tag{2.26}$$

$$\phi(q)\phi(q^3) = a(q^4) + 2q\psi(q^2)\psi(q^6), \tag{2.27}$$

$$a(q) = a(q^4) + 6q\psi(q^2)\psi(q^6), \tag{2.28}$$

$$\psi(q)\psi(q^3) = \psi(q^4)\phi(q^6) + q\phi(q^2)\psi(q^{12}), \tag{2.29}$$

$$4\phi(q^3)\phi(q)a(q^2)-\phi(q)^4=3\phi(q^3)^4, \tag{2.30}$$

$$c(q)=3\frac{E^3(q^3)}{E(q)}. \tag{2.31}$$

At this point, it is expedient to introduce a projection operator $P_{t,s}$. It is defined by its action on power series as follows. Let t,s be non-negative integers such that $0\leq s<t$. Then

$$P_{t,s}\sum_{n\geq 0}c(n)q^n:=\sum_{n\geq 0}c(tn+s)q^{tn+s}. \tag{2.32}$$

Next, we observe that

$$P_{2,1}f(q^3,q^{15})\phi(q^3)=\sum_{x\not\equiv y \pmod 2} q^{(3x+1)^2+3y^2-1}=\sum_{x,y} q^{(3x+1)^2+3(x+2y+1)^2-1}=q^3c(q^{12}). \tag{2.33}$$

Replacing q^3 by q in (2.33), we find

$$P_{2,1}f(q,q^5)\phi(q)=qc(q^4). \tag{2.34}$$

We note in passing, that (2.34) follows easily from

$$f(q,q^5)\phi(q)=qc(q^4)+\psi(q^2)f(q^2,q^4), \tag{2.35}$$

but we will not take the time to prove it. A motivated reader is invited to examine [1] and [13] for many similar relations.

With the aid of (2.21) it is straightforward to verify that

$$P_{3,1}\phi(q)^2\phi(q^3)=4qf(q^3,q^{15})\phi(q^3)\phi(q^9). \tag{2.36}$$

Next, we employ (2.22), (2.27), (2.28), (2.29), (2.34) along with (2.36) to deduce

$$P_{24,1}\phi(q)^2\phi(q^3)=4qf(q^{24},q^{48})a(q^{48})+8q^{25}\psi(q^{72})c(q^{48}). \tag{2.37}$$

We would like to transform (2.37) into

$$P_{24,1}\phi(q)^2\phi(q^3)=4q\frac{E(q^{24})^5}{E(q^{48})^2}+16q^{25}\psi(q^{72})c(q^{48}). \tag{2.38}$$

This equation will play an important role in establishing Theorem 1.11.

Proof. Comparing (2.37) and (2.38) we see that all that is needed to show is

$$a(q^2)=\frac{E(q)^5}{f(q,q^2)E(q^2)^2}+2q\psi(q^3)\frac{c(q^2)}{f(q,q^2)}. \tag{2.39}$$

Replacing q by $-q$ in (2.39) and multiplying both sides by $-4\phi(q^3)\phi(q)$ with the aid of (2.3), (2.9), (2.12), (2.13), (2.14) and (2.31), we get

$$3\phi(q)^4-(4\phi(q^3)\phi(q)a(q^2)-\phi(q)^4)=24qf(q,q^5)^3\phi(q^3). \tag{2.40}$$

Making use of (2.30) we arrive at (2.26). The proof of (2.38) is now complete. □

3. Proof of Theorems 1.9, 1.10 and 1.11

We begin by rewriting Theorem 1.9 as

Theorem 3.1. *Let n be a positive integer. Then,*

$$[q^{8n+1}]q^9\psi(q^8)\psi(q^{32})^2 \geq 0, \tag{3.1}$$

with equality if and only if $8n+1=M^2$ and all prime divisors of M are congruent to $1 \pmod 4$.

Proof. Replace q by $-q$ in (2.25) to get

$$\phi(-q)^2 = \phi(q^2)^2 - 4q\psi(q^4)^2. \tag{3.2}$$

Subtracting (3.2) from (2.25) and replacing q by q^8 we find

$$\phi(q^8)^2 - \phi(-q^8)^2 = 8q^8\psi(q^{32})^2. \tag{3.3}$$

Hence

$$\phi(q^8)^2 - 8q^8\psi(q^{32})^2 = \phi(-q^8)^2. \tag{3.4}$$

Multiplying both sides of (3.4) by $q\psi(q^8)$ and employing (2.15) we obtain

$$q\phi(q^8)^2\psi(q^8) - qE(q^8)^3 = 8q^9\psi(q^{32})^2\psi(q^8) \tag{3.5}$$

It is straightforward to verify that

$$P_{4,1}\frac{\phi(q)^3}{6} = q\phi(q^4)^2\psi(q^8). \tag{3.6}$$

With the aid of (2.25) and (3.6) we obtain

$$P_{8,1}\frac{\phi(q)^3}{6} = q\phi(q^8)^2\psi(q^8), \tag{3.7}$$

$$P_{8,5}\frac{\phi(q)^3}{24} = q^5\psi(q^{16})^2\psi(q^8). \tag{3.8}$$

Combining (2.7) and (3.5), (3.7) we find that

$$P_{8,1}\left(\frac{\phi(q)^3}{6} - \sum_{n>0}\left(\frac{-4}{n}\right)nq^{n^2}\right) = 8q^9\psi(q^{32})^2\psi(q^8). \tag{3.9}$$

Therefore, if $8n+1$ is not a perfect square, then

$$[q^{8n+1}]q^9\psi(q^8)\psi(q^{32})^2 = \frac{(1,1,1,0,0,0;8n+1)}{48} > 0, \tag{3.10}$$

by (1.2). If $8n+1=M^2$, with M being a positive integer, then

$$\left(\frac{-4}{M}\right) = \left(\frac{-1}{M}\right) \tag{3.11}$$

and

$$[q^{M^2}]8q^9\psi(q^8)\psi(q^{32})^2 = \frac{(1,1,1,0,0,0;M^2)}{6} - \left(\frac{-1}{M}\right)M \geq 0, \tag{3.12}$$

with equality if and only if all prime divisors of M are congruent to $1 \pmod 4$, by (1.16). □

Next, we turn to Theorem 1.10. We rewrite it as

Theorem 3.2. *Let n be a positive integer. Then,*

$$[q^{8n+1}]q^{17}\phi(q^8)\psi(q^8)\psi(q^{128}) \geq 0, \tag{3.13}$$

with equality if and only if $8n+1=E^2$ and all prime divisors of E are congruent to 1 or 3 modulo 8.

Proof. Replace q by $-q$ in (2.20) to get

$$\phi(-q)=\phi(q^4)-2q\psi(q^8). \tag{3.14}$$

Subtracting (2.20) from (3.14) and replacing q by q^2 we find

$$\phi(-q^2)=\phi(q^2)-4q^2\psi(q^{16}). \tag{3.15}$$

Multiply both sides (3.15) by $\phi(q)\psi(q)$ and use (2.16) on the left to get

$$\phi(q)^2\psi(-q)=\phi(q^2)\phi(q)\psi(q)-4q^2\psi(q^{16})\phi(q)\psi(q). \tag{3.16}$$

Replacing q by q^8 in (3.16) and multiplying both sides by q we obtain

$$q\phi(q^8)^2\psi(-q^8)=q\phi(q^{16})\psi(q^8)\phi(q^8)-4q^{17}\phi(q^8)\psi(q^8)\psi(q^{128}). \tag{3.17}$$

Clearly (2.15) implies that

$$q\phi(q^8)^2\psi(-q^8)=qE(-q^8)^3, \tag{3.18}$$

and so we established that

$$q\psi(q^8)\phi(q^8)\phi(q^{16})-qE(-q^8)^3=4q^{17}\phi(q^8)\psi(q^8)\psi(q^{128}) \tag{3.19}$$

holds true. It is straightforward to verify that

$$P_{8,1}\frac{\phi(q)^2\phi(q^2)}{4}=q\psi(q^8)\phi(q^8)\phi(q^{16}). \tag{3.20}$$

Combining (2.8) and (3.19), (3.20) we find that

$$P_{8,1}\left(\frac{\phi(q)^2\phi(q^2)}{4}-\sum_{n>0}\left(\frac{-2}{n}\right)nq^{n^2}\right)=4q^{17}\phi(q^8)\psi(q^8)\psi(q^{128}). \tag{3.21}$$

And so, if $8n+1$ is not a perfect square, then

$$[q^{8n+1}]q^{17}\phi(q^8)\psi(q^8)\psi(q^{128})=\frac{(1,1,2,0,0,0;8n+1)}{16}>0, \tag{3.22}$$

by (1.6). If $8n+1=E^2$, with E being positive integer, then

$$[q^{E^2}]4q^{17}\phi(q^8)\psi(q^8)\psi(q^{128})=\frac{(1,1,2,0,0,0;E^2)}{4}-E\left(\frac{-2}{E}\right)\geq 0, \tag{3.23}$$

with equality if and only if all prime divisors of E are congruent to 1 or 3 modulo 8, by (1.17). □

It remains to prove Theorem 1.11. We begin by rewriting it as

Theorem 3.3. *Let n be a positive integer. Then,*

$$[q^{24n+1}]q^{25}\psi(q^{72})c(q^{48})\geq 0, \tag{3.24}$$

with equality if and only if $24n+1=W^2$ and all prime divisors of W are congruent to 1 modulo 3.

Proof. From (2.18) and (2.38) we have

$$P_{24,1}\left(\phi(q)^2\phi(q^3)-4\sum_{n>0}\left(\frac{-12}{n}\right)nq^{n^2}\right)=16q^{25}\psi(q^{72})c(q^{48}). \tag{3.25}$$

If $24n+1$ is not a perfect square, then

$$[q^{24n+1}]q^{25}\psi(q^{72})c(q^{48})=\frac{(1,1,3,0,0,0;24n+1)}{16}>0, \tag{3.26}$$

by (1.7). If $24n+1=W^2$, with W being positive integer, then

$$[q^{W^2}]4q^{25}\psi(q^{72})c(q^{48})=\frac{(1,1,3,0,0,0;W^2)}{4}-W\left(\frac{-3}{W}\right)\geq 0, \tag{3.27}$$

with equality if and only if all prime divisors of W are congruent to 1 modulo 3, by (1.18). □

4. On spinor-regular ternary forms of discriminant 16384

According to Dickson [10], a positive definite ternary quadratic form is said to be regular if it represents all natural integers not excluded by congruences; that is, it represents all integers represented by its genus. We now know that there are exactly 913 regular ternary quadratic forms [16]. In [17], Jones and Pall discussed genus companions of certain regular diagonal forms. These companions are almost regular, but they fail to represent an infinite set of natural numbers in the same square-class. For example, ternary quadratic form $(3,4,9,0,0,0)$ happens to be almost regular. It is the only genus mate of regular form $(1,3,36,0,0,0)$. To demonstrate the usefulness of a special case of the quintuple product identity (2.19), we digress a little by taking the time to prove the following

Lemma 4.1. *Let n be a non-negative integer. Then*

$$(3,4,9,0,0,0;n) \geq (1,3,36,0,0,0;n), \tag{4.1}$$

provided n is not of the form $(6j+r)^2$ with $j \in \mathbb{N}$ and $r = 1, 2$. Moreover,

$$(1,3,36,0,0,0;(6j+r)^2) > (3,4,9,0,0,0;(6j+r)^2), \tag{4.2}$$

$$(3,4,9,0,0,0;(6j+r)^2) \geq 0, \tag{4.3}$$

with equality if and only if all prime divisors of $6j+r$ are congruent to 1 modulo 3.

Proof. We begin by setting $s = 1, m = 12$ in [Thm. 2.1, [5]] to get

$$\sum_{x,y} q^{72x^2+12xy+y^2} - \sum_{x,y} q^{72x^2+60xy+13y^2} = 2qE(q^{12})^2. \tag{4.4}$$

The unimodular transformation $x \mapsto y$, $y \mapsto -x-6y$ yields

$$\sum_{x,y} q^{72x^2+12xy+y^2} = \phi(q)\phi(q^{36}). \tag{4.5}$$

Analogously, with the aid of $x \mapsto -x+y$, $y \mapsto 2x-3y$ we find that

$$\sum_{x,y} q^{72x^2+60xy+13y^2} = \phi(q^4)\phi(q^9). \tag{4.6}$$

Hence

$$\vartheta(1,3,36,0,0,0,q) - \vartheta(3,4,9,0,0,0,q) = 2q\phi(q^3)E(q^{12})^2. \tag{4.7}$$

From (2.19) and (4.7) it follows that

$$(3,4,9,0,0,0;n) = (1,3,36,0,0,0;n), \tag{4.8}$$

if n is not a perfect square, and that for $m \in \mathbb{N}$

$$(1,3,36,0,0,0;m^2) - (3,4,9,0,0,0;m^2) = 2(-1)^{m+1}\left(\frac{-3}{m}\right)m. \tag{4.9}$$

Hence

$$(3,4,9,0,0,0;9m^2) = (1,3,36,0,0,0;9m^2). \tag{4.10}$$

Finally, (1.19) implies that for $j \in \mathbb{N}$

$$(3,4,9,0,0,0;(6j+4)^2) > (1,3,36,0,0,0;(6j+4)^2) > 0, \tag{4.11}$$

$$(3,4,9,0,0,0;(6j+5)^2) > (1,3,36,0,0,0;(6j+5)^2) > 0, \tag{4.12}$$

$$(1,3,36,0,0,0;(6j+2)^2) > (3,4,9,0,0,0;(6j+2)^2) > 0, \tag{4.13}$$

$$(1,3,36,0,0,0;(6j+1)^2) > (3,4,9,0,0,0;(6j+1)^2), \tag{4.14}$$

and

$$(3,4,9,0,0,0;(6j+1)^2) \geq 0, \tag{4.15}$$

with equality if and only if all prime divisors of $(6j+1)$ are congruent to 1 modulo 3. □

Hence $3x^2+4y^2+9z^2$ represents exclusively all positive integers represented by $x^2+3y^2+36z^2$, which are not of the form W^2, where W is generated by 1 and primes congruent to 1 modulo 3. Recalling [10, Table 5], we have

Theorem 4.2. *The form $3x^2+4y^2+9z^2$ represents exclusively all positive integers not of the form*
$(3m+2)$,
$(4m+2)$,
$9^a(9m+6)$,
W^2,
where $a,m,W\in\mathbb{N}$ and W is generated by 1 and primes congruent to 1 modulo 3.

Nowadays, the Jones-Pall forms are called spinor regular. The list containing 29 spinor regular forms was put together in [15]. It contains seven original Jones-Pall forms along with eight forms from [2] and ten forms from [18]. Jagy contributed four forms to this list. In addition, he checked all positive ternary form with discriminant up to $1,400,000$ looking for all the spinor regular forms, based on a recipe in [8]. The list of 29 forms is conjectured to be complete.

All 10 of Kaplansky's forms in [18] can be easily handled by the methods discussed in this paper. Here, we focus on just two forms $(9,16,36,16,4,8)$ and $(9,17,32,-8,8,16)$, both of discriminant 16384. We begin by proving the following

Lemma 4.3.

$$\vartheta(9,16,36,16,4,8,q)=\phi(q^{64})^3+2q^{16}\psi(q^{128})\phi(q^{64})^2+8q^{36}\psi(q^{32})\psi(q^{128})^2+2q^9\psi(q^8)\psi(q^{32})^2. \tag{4.16}$$

Proof. It is easy to check that

$$9x^2+16y^2+36z^2+16yz+4xz+8xy=(x+4y+2z)^2+8x^2+32z^2. \tag{4.17}$$

Hence

$$\vartheta(9,16,36,16,4,8,q)=\sum_{x,y,z} q^{(2x+4y+2z)^2+8(2x)^2+32z^2}+\sum_{x,y,z} q^{(2x+1+4y+2z)^2+8(2x+1)^2+32z^2}. \tag{4.18}$$

Next,

$$\begin{aligned}\sum_{x,y,z} q^{(2x+4y+2z)^2+8(2x)^2+32z^2} &= \sum_{x\equiv z \pmod 2,\atop y} q^{(2x+4y+2z)^2+32x^2+32z^2}\\ &+\sum_{x\not\equiv z \pmod 2,\atop y} q^{(2x+4y+2z)^2+32x^2+32z^2}.\end{aligned} \tag{4.19}$$

Using the substitutions $x\mapsto x+z$, $z\mapsto x-z$, $y\mapsto y-x$ and $x\mapsto x+z+1$, $z\mapsto x-z$, $y\mapsto y-x$ we find that

$$\begin{aligned}\sum_{x\equiv z \pmod 2,\atop y} q^{(2x+4y+2z)^2+32x^2+32z^2} &= \sum_{x,y,z} q^{16y^2+64x^2+64z^2}\\ &= \sum_{x,y,z} q^{64y^2+64x^2+64z^2}+\sum_{x,y,z} q^{16(2y+1)^2+64x^2+64z^2}\\ &= \phi(q^{64})^3+2q^{16}\psi(q^{128})\phi(q^{64})^2,\end{aligned} \tag{4.20}$$

and

$$\begin{aligned}\sum_{x\not\equiv z \pmod 2,\atop y} q^{(2x+4y+2z)^2+32x^2+32z^2} &= \sum_{x,y,z} q^{(4y+2)^2+16(2x+1)^2+16(2z+1)^2}\\ &= 8q^{36}\psi(q^{32})\psi(q^{128})^2,\end{aligned} \tag{4.21}$$

respectively.
Hence

$$\sum_{x,y,z} q^{(2x+4y+2z)^2+8(2x)^2+32z^2} = \phi(q^{64})^3 + 2q^{16}\psi(q^{128})\phi(q^{64})^2 + 8q^{36}\psi(q^{32})\psi(q^{128})^2. \tag{4.22}$$

Analogously,

$$\begin{aligned}\sum_{x,y,z} q^{(2x+1+4y+2z)^2+8(2x+1)^2+32z^2} &= \sum_{x\equiv z \pmod 2,\atop y} q^{(2x+1+4y+2z)^2+8(2x+1)^2+32z^2}\\ &+ \sum_{x\not\equiv z \pmod 2,\atop y} q^{(2x+1+4y+2z)^2+8(2x+1)^2+32z^2}\\ &= \sum_{x,y,z} q^{(4y+1)^2+4(4x+1)^2+4(4z+1)^2}\\ &+ \sum_{x,y,z} q^{(4y+3)^2+4(4x+3)^2+4(4z+3)^2}\\ &= 2q^9\psi(q^8)\psi(q^{32})^2.\end{aligned} \tag{4.23}$$

Combining (4.18), (4.22) and (4.23), we arrive at (4.16). □

Next, employing (4.16) together with (3.6), we have the following

Lemma 4.4. *Let n be a non-negative integer. We have*

$$(9,16,36,16,4,8;64n) = (1,1,1,0,0,0;n) \geq 0, \tag{4.24}$$

with equality if and only if n is of the form $4^a(8m+7)$ with $a, m \in \mathbb{N}$,

$$(9,16,36,16,4,8;64n+16) = \frac{(1,1,1,0,0,0;4n+1)}{3} > 0, \tag{4.25}$$

$$(9,16,36,16,4,8;32n+4) = 8[q^{32n+4}]q^{36}\psi(q^{32})\psi(q^{128})^2 \geq 0, \tag{4.26}$$

$$(9,16,36,16,4,8;8n+1) = 2[q^{8n+1}]q^9\psi(q^8)\psi(q^{32})^2 \geq 0, \tag{4.27}$$

$$(9,16,36,16,4,8;8n+r) = 0, \quad \text{if } r \in \{3,5,7\}, \tag{4.28}$$

$$(9,16,36,16,4,8;64n+2\tilde{r}) = 0, \quad \text{if } \tilde{r} \notin \{0,2,8,18\}. \tag{4.29}$$

Making use of (3.1) with $q \mapsto q$, $q \mapsto q^4$, we establish

Theorem 4.5. *The form $9x^2+16y^2+36z^2+16yz+4xz+8xy$ represents exclusively all positive integers not of the form*
$4^a(8m+7)$,
$4^a(8m+3), 0 \leq a \leq 2$,
$4^a(4m+2), 0 \leq a \leq 2$,
$4^a(8m+5), 0 \leq a \leq 1$,

$M^2, 4M^2$,
where $a, m, M \in \mathbb{N}$ *and* M *is generated by* 1 *and primes congruent to* 1 *modulo* 4.

We now turn to the second form $(9, 17, 32, -8, 8, 16)$. Again, we start with the easily verifiable identity

$$9x^2 + 17y^2 + 32z^2 - 8yz + 8xz + 6xy = (x + 3y - 4z)^2 + 4(x - y)^2 + 4(x + y + 2z)^2, \tag{4.30}$$

after a bit of labor, to obtain

$$\begin{aligned}\vartheta(9, 17, 32, -8, 8, 6, q) = \phi(q^{32})^2\phi(q^{256}) + 2q^{20}\psi(q^{32})\psi(q^{64})^2 + 8q^{80}\psi(q^{128})\psi(q^{256})^2 \\ + 16q^{144}\psi(q^{128})\psi(q^{512})^2 + 4q^{36}\psi(q^{32})\psi(q^{128})^2 + 2q^9\psi(q^8)\psi(q^{32})^2.\end{aligned} \tag{4.31}$$

Hence, recalling (3.8), we have the following

Lemma 4.6. *Let* $n, m, a \in \mathbb{N}$. *We have*

$$(9, 17, 32, -8, 8, 6; 32n) = (1, 1, 8, 0, 0, 0; n) \geq 0, \tag{4.32}$$

with equality if and only if n *is of the form* $2\ 4^a(8m + 7), 4m + 3$ *or* $2(8m + 3)$ *by* [10, Table 5]

$$(9, 17, 32, -8, 8, 6; 32n + 20) = \frac{(1, 1, 1, 0, 0, 0; 8n + 5)}{12} > 0, \tag{4.33}$$

$$(9, 17, 32, -8, 8, 6; 128n + 80) = \frac{(1, 1, 1, 0, 0, 0; 8n + 5)}{3} > 0, \tag{4.34}$$

$$(9, 17, 32, -8, 8, 6; 128n + 16) = 16[q^{128n+16}]q^{144}\psi(q^{128})\psi(q^{512})^2 \geq 0, \tag{4.35}$$

$$(9, 17, 32, -8, 8, 6; 32n + 4) = 4[q^{32n+4}]q^{36}\psi(q^{32})\psi(q^{128})^2 \geq 0, \tag{4.36}$$

$$(9, 17, 32, -8, 8, 6; 8n + 1) = 2[q^{8n+1}]q^9\psi(q^8)\psi(q^{32})^2 \geq 0, \tag{4.37}$$

$$(9, 17, 32, -8, 8, 6; 8n + r) = 0, \quad \text{if } r \in \{3, 5, 7\}, \tag{4.38}$$

$$(9, 17, 32, -8, 8, 6; 64n + 2\tilde{r}) = 0, \quad \text{if } \tilde{r} \notin \{0, 2, 8, 16, 18\}. \tag{4.39}$$

Employing (3.1) with $q \mapsto q$, $q \mapsto q^4$ and $q \mapsto q^{16}$, we arrive at

Theorem 4.7. *The form* $9x^2 + 17y^2 + 32z^2 - 8yz + 8xz + 6xy$ *represents exclusively all positive integers not of the form*
$4^a(8m + 7)$,
$4^a(8m + 6), 0 \leq a \leq 2$,
$4^a(8m + 3), 0 \leq a \leq 3$,
$4^a(8m + 2), 0 \leq a \leq 1$,
$8m + 5$,
$M^2, 4M^2, 16M^2$,
where $a, m, M \in \mathbb{N}$ *and* M *is generated by* 1 *and primes congruent to* 1 *modulo* 4.

5. Acknowledgements

I am grateful to William Jagy for a most illuminating discussion and to Shaun Cooper, Michael Hirschhorn, Elizabeth Loew, Frank Patane for a careful reading of the manuscript. I would like to thank George Andrews, who encouraged me to write up these observations.

References

[1] P. Barrucand, S. Cooper and M. D. Hirschhorn, *Results of Hurwitz type for five or more squares*, Ramanujan J.,**6**, (2002), 347–367.

[2] J. W. Benham, A. G. Earnest, J. S. Hsia, D. C. Hung, *Spinor regular Positive Ternary Quadratic Forms*, J. London Math. Soc. **42**, (1990), 1–10.

[3] A. Berkovich, *On representation of an integer by* $X^2 + Y^2 + Z^2$ *and the modular equations of degree* 3 *and* 5, Quadratic and higher degree forms, 29–49, DEVM, **31**, Springer, New York, 2013.

[4] A. Berkovich, W. C. Jagy, *On representation of an integer as the sum of three squares and the ternary quadratic forms with the discriminants* p^2*,* $16p^2$, J. Number Theory, **132**, (2012), 259-274.

[5] A. Berkovich, F. Patane, *Binary quadratic forms and the Fourier coefficients of certain weight* 1 *eta-quotients*, J. Number Theory **135**, (2014), 185-220.

[6] B. C. Berndt, *Ramanujan's Notebooks, Part III*, Springer, New York, 1991.

[7] B. C. Berndt, *Number Theory in the Spirit of Ramanujan*, Student Mathematical Library, **34** AMS, Providence, RI, 2006.

[8] W. K. Chan, A. G. Earnest, *Discriminant bounds for spinor regular ternary quadratic lattices*, J. London Math. Soc. **69**, (2004), 545-561.

[9] S. Cooper, H. Y. Lam, *On the Diophantine equation* $n^2 = x^2 + by^2 + cz^2$, J. Number Theory, **133**, (2013), 719-737.

[10] L. E. Dickson, *Modern Elementary Theory of Numbers*, Univ. of Chicago Press, 1939.

[11] C. F. Gauss, *Disquisitiones Arithmeticae*, Yale University Press, 1965.

[12] X. Guo, Y. Peng, H. Qin, *On the representation numbers of ternary quadratic forms and modular forms of weight* 3/2, J. Number Theory, **140**, (2014), 235-266.

[13] M. D. Hirschhorn, F. G. Garvan, J. M. Borwein, *Cubic analogues of the Jacobian theta function* $\theta(z, q)$, Canad. J. Math. **45** (1993), no. 4, 673–694.

[14] A. Hurwitz, *Mathematische Werke*, Band2, 751, Birkhauser, Basel, 1933.

[15] W. C. Jagy, *Integral Positive Ternary Quadratic Forms*, arXiv:1010.3677v1 [math.NT].

[16] W. C. Jagy, I. Kaplansky, A. Schiemann, *There are* 913 *regular ternary quadratic forms*, Mathematika, **44** (1997), 332–341.

[17] B. W. Jones, G. Pall, *Regular and semi-regular positive ternary quadratic forms*, Acta Mathematica **70**(1939), 165–191.

[18] I. Kaplansky, *Notes on Ternary Forms, III, JP (Jones-Pall) forms*, Unpublished, 1996, three pages.

[19] A. M. Legendre, *Essai sur la theorie des nombres*, Cambridge University Press, 2009.

[20] C. D. Olds, *On the representations,* $N_3(n^2)$, Bull. Amer. Math. Soc. **47**, (1941), 499–503.

[21] G. Pall, *On the arithmetic of quaternions*,Trans. Amer. Math. Soc. **47** (1940), 487–500.

[22] L. C. Shen, *On the modular equations of degree* 3, Proc. Amer. Math. Soc. **122** (1994), no. 4, 1101–1114.

[23] C. L. Siegel, *Lectures on the Analytical Theory of Quadratic Forms*, Notes by Morgan Ward. Third revised edition. Buchhandlung Robert Peppmüller, Göttingen, 1963.

Department of Mathematics, University of Florida, 496 Little Hall, Gainesville, FL 32611, USA
E-mail address: alexb@ufl.edu